

## HEDIS® Provider Guide:

### Measure Description

Women 16-24 years of age who were identified as sexually active and who had at least one Chlamydia test during the measurement year.

Members identified as sexually active:

- Pregnancy
- STI screening
- Contraceptive prescribed
- Pregnancy test completed

### Using Correct Billing Codes

Description	CPT Code
Chlamydia Screening	87110, 87270, 87320, 87490-87492, 87810

Contraceptives	
<ul style="list-style-type: none"> <li>• Desogestrel-ethinyl estradiol</li> <li>• Dienogest-estradiol (multiphasic)</li> <li>• Drospirenone-ethinyl estradiol</li> <li>• Drospirenone-ethinyl estradiol-levomefolate (biphasic)</li> <li>• Ethinyl estradiol-ethynodiol</li> <li>• Ethinyl estradiol-etonogestrel</li> <li>• Ethinyl estradiol-levonorgestrel</li> <li>• Ethinyl estradiol-norelgestromin</li> </ul>	<ul style="list-style-type: none"> <li>• Ethinyl estradiol-norethindrone</li> <li>• Ethinyl estradiol-norgestimate</li> <li>• Ethinyl estradiol-norgestrel</li> <li>• Etonogestrel</li> <li>• Levonorgestrel</li> <li>• Medroxyprogesterone</li> <li>• Mestranol-norethindrone</li> <li>• • Norethindrone</li> </ul>

### How to Improve HEDIS® Scores

- Perform Chlamydia screening every year on every 16-24-year old female identified as sexually active (use any visit opportunity).
- Add Chlamydia screening as a standard lab for women 16-24 years old. Use adolescent well-care exams and well woman exams for this purpose.
- Ensure that you have an opportunity to speak with your adolescent female patient without her parents present.
- Remember that Chlamydia screening can be performed through a urine test. Offer this option for your patients.